

Leaf gas exchange responses of 13 prairie grassland species to elevated CO₂ and increased nitrogen supply

Tali D. Lee¹, Mark G. Tjoelker², David S. Ellsworth³ and Peter B. Reich¹

¹Department of Forest Resources, University of Minnesota, 1530 Cleveland Ave. N., St. Paul, MN 55108, USA; ²Department of Forest Science, Texas A & M University, College Station, TX, 77843–2135, USA; ³Environmental Sciences Department, Brookhaven National Laboratory, Upton, NY 11973, USA

Summary

Author for correspondence:

Tali D. Lee

Tel: +1 612 624 3671

Fax: +1 612 6255212

Email: tlee@biosci.cbs.umn.edu

Received: 11 September 2000

Accepted: 12 January 2001

- Leaf gas exchange responses to elevated CO₂ and N are presented for 13 perennial species, representing four functional groups: C₃ grasses, C₄ grasses, legumes, and nonleguminous forbs. Understanding how CO₂ and N effects interact is important to predict plant community response to global change.
- Plants were field-grown in monoculture under current ambient and elevated (560 μmol mol⁻¹) CO₂ concentrations (free-air CO₂ enrichment), in combination with soil N treatments, for two growing seasons.
- All species, regardless of functional group, showed pronounced photosynthetic acclimation to elevated CO₂, resulting in minimal stimulation of photosynthesis (A) averaging +15% in C₃ grasses, +8% in forbs, +7% in legumes and –2% in C₄ grasses. The effects of CO₂ and soil N supply did not interact for any leaf traits measured. Elevated CO₂ consistently decreased stomatal conductance (g_s) leading to 40% increase in A/g_s.
- This substantial acclimation of photosynthesis was greater in magnitude than in most field studies, and was associated with the combined effects of decreased g_s and decreased leaf N concentrations in response to growth under elevated CO₂.

Key words: elevated CO₂, nitrogen availability, photosynthetic acclimation, functional groups, prairie grassland, stomatal conductance.

© *New Phytologist* (2001) **150**: 000–000

Introduction

Rising atmospheric carbon dioxide (CO₂) concentration is predicted to have profound effects on ecosystems. Human activity is also altering the global nitrogen (N) cycle by substantially increasing the inputs of fixed forms of N, primarily by the extensive use of chemical fertilizers and combustion of fossil fuels (Vitousek *et al.*, 1997). The effects of elevated atmospheric CO₂ and increased soil N on vegetation likely interact in complex ways and differently at different scales. Although larger-scale field experiments are increasing in number and longevity, there continues to be a need for field experiments to determine the extent to which results found in controlled environment studies apply to intact plant communities (Polley, 1997).

Plant responses to elevated CO₂ are fundamentally mediated by photosynthesis (Drake *et al.*, 1997), and can potentially lead to a suite of morphological and growth changes. It is well documented that increased CO₂ enhances the photosynthetic rate and growth of most C₃ plants (Bowes, 1993). However,

as larger-scale, longer-term studies are being conducted, findings indicate the degree of this response to be variable and its persistence over the long-term questionable. To date, photosynthetic responses to experimentally doubled CO₂ levels have ranged from neutral, even negative, to strongly positive in most crop systems where typical increases average from 20 to 40% (Schimel, 1995). Responses of less studied wild species in natural systems are often considerably lower in magnitude than crops and in some cases under protracted exposure to elevated CO₂, photosynthetic rates decline, resulting in a complete lack of enhancement (Bowes, 1993; Wand *et al.*, 1999). A reduction in photosynthetic capacity with exposure to elevated CO₂ may occur (e.g. photosynthetic acclimation; Gunderson & Wullschlegel, 1994), in connection with changes in leaf chemistry and structure, as well as feedbacks governed by whole plant growth dynamics. Common responses to CO₂ enrichment include: decreases in the amount or activity of Rubisco, increases in total nonstructural carbohydrate concentrations, and decreases in leaf N concentration and leaf area to mass ratios (Curtis, 1996), which collectively should

lead to decreased rates of photosynthesis (Reich *et al.*, 1997). Understanding acclimation of photosynthesis to increased atmospheric CO₂ concentration, and how soil N availability affects this response, is critical for predicting plant community responses to environmental change.

Potential limitations of resources such as nutrients and light may mitigate photosynthetic responses to elevated CO₂ (Drake *et al.*, 1997). Considering the dynamics and influence of N on photosynthesis and growth becomes critical in the attempt to characterize individual plant, as well as ecosystem, responses to elevated CO₂ (Vitousek *et al.*, 1997). Since biologically available N currently limits productivity in most ecosystems, and because tissue N is a major determinant of photosynthesis (Nijs *et al.*, 1995; Reich *et al.*, 1997), low N may limit potential photosynthetic enhancement under elevated CO₂. Several simulation models predict that plant CO₂ responses are constrained by N limitation (McMurtie & Comins, 1996; Rastetter *et al.*, 1997), although actual evidence is mixed (Poorter, 1998), ranging from no consistent effect of nutrient availability on plant responsiveness to elevated CO₂ (Idso & Idso, 1994; Lloyd & Farquhar, 1996) to a decreased CO₂ sensitivity which is linked to low nutrient availability (Larigauderie *et al.*, 1988; Bazzaz, 1990; Oechel *et al.*, 1994; Leadley & Körner, 1996; Poorter, 1998). Thus, a plant's response to limiting factors other than atmospheric CO₂ may have a great impact on how it responds to elevated CO₂ (Bowes, 1993).

Research on differential responses to CO₂ among plant functional groups may help explain the large degree of interspecific variation in acclimation to elevated CO₂. As a way to model the response of complex ecosystems based on group rather than species parameterizations, functional groups have been proposed because they have been found to help explain variations in species responses to the environment (Díaz, 1995). However, Díaz (1995) points out that plant responsiveness to elevated CO₂ may involve traits not usually considered in functional group definitions. Evaluating species in more natural field settings provides the opportunity to identify general patterns associated with strong or weak responses to elevated CO₂ and to test the usefulness of these functional classifications.

The overall objective of this experiment was to investigate how elevated CO₂ concentrations and increased soil N interact to affect leaf-level physiological processes of a variety of wild perennial plant species in a field setting and to examine if these responses to elevated CO₂ and N remain similar across two growing seasons. Our evaluation was done at the leaf level in order to identify potential physiological mechanisms underlying ecosystem response to these global change elements. In this study, the following questions are addressed: to what extent do field grown prairie species acclimate leaf photosynthesis to elevated CO₂ concentrations and is this response modulated by soil N supply?; and do functional groupings help explain the variation in species responses to elevated CO₂ and increased soil N?

Materials and Methods

Research site

The study site is located at the Cedar Creek Natural History Area in east central Minnesota, USA (Lat. 45° N, Long. 93° W). The soils are sandy, derived from a glacial outwash sandplain, and previous experiments have determined nitrogen to be the major soil resource that limits plant growth (Tilman, 1987). N mineralization rates for grassland soils at Cedar Creek are estimated to range between 2 and 3 g m⁻² yr⁻¹ (Wedin & Tilman, 1996). Cedar Creek has a continental climate with cold winters (mean January temperature = -11°C), warm summers (mean July temperature = 22°C), and mean annual precipitation totaling 660 mm yr⁻¹. The average maximum daily temperature and total precipitation for the 1998 and 1999 growing seasons (April–September) was 25°C with 389 mm rainfall in 1998 and 24°C with 637 mm rainfall in 1999.

Experimental design and the FACE system

The overall experiment, referred to as BioCON (Biodiversity, Carbon dioxide, and Nitrogen effects on ecosystem functioning, <http://www.swan.lter.umn.edu/biocon/>), was established in 1997 on secondary successional grassland after removing previous vegetation (Reich *et al.*, 2001a,b). The study site consisted of six circular areas (20-m diameter), each containing 61–2 × 2 m plots. The experimental treatments were arranged in factorial combination of CO₂ concentration (368 or 560 µmol mol⁻¹) and soil N supply (low or high (4 g N m⁻² yr⁻¹ added)) with each species in monoculture replicated twice for every CO₂ × N level. The design consisted of a split-plot arrangement of treatments in a randomized design with CO₂ treatment as the whole-plot factor, which was replicated three times among the six rings. The subplot factor of soil N treatment was randomly assigned to individual plots among the six rings. CO₂ was applied using free-air CO₂ enrichment (FACE) technology (Lewin *et al.*, 1994) during all daylight hours during the growing season from April 9 to October 16, 1998 and from April 20 to November 9, 1999. One-minute averages were within 10% of the target concentration 94% and 95% of the time in 1998 and 1999, respectively. The high N plots were amended with 4 g N m⁻² yr⁻¹, as ammonium nitrate (NH₄NO₃) in solid form, in May, June and July of each year. Monoculture plots of 13 species, representing four functional classifications of plants based on similarities in physiology and growth forms: C₃ grasses, C₄ grasses, legumes, and nonleguminous forbs, were chosen for leaf-level physiological measurements in this study. These species include: C₃ grasses: *Agropyron repens* (L.) Beauv., *Bromus inermis* Leyss., *Koeleria cristata* Pers.; C₄ grasses: *Andropogon gerardii* Vit., *Schizachyrium scoparium* (Mich.) Nash, *Sorghastrum nutans* (L.) Nash; nonleguminous forbs: *Achillea millefolium* L.,

Anemone cylindrica A. Gray, *Solidago rigida* L.; and legumes: *Lupinus perennis* L., *Lespedeza capitata* Mich., *Amorpha canescens* Pursh, *Petalostemum villosum* Nutt. Species hereafter are referred to by their genus. Most species were measured in both 1998 and 1999, however, due to low biomass *Lespedeza* was measured only in 1998 and *Amorpha* and *Petalostemum* were measured only in 1999. For plot biomass data refer to Reich *et al.* (2001).

Gas exchange and leaf nitrogen

1 During both the 1998 and 1999 growing seasons, *in situ* rates of leaf net photosynthesis (A) were measured using CIRAS-1 portable infrared gas exchange systems (PP Systems, Hitchin, UK) operated in open-configuration with controlled temperature, CO₂ concentration, and vapor pressure. Measurements were made on an upper fully expanded leaf of an individual plant representing each monoculture plot, typically between 09:00 and 15:00 hours local time. Since leaf traits vary with age, all measures were made using leaves of similar ontogenetic stage. We used upper fully expanded young to mid-aged leaves, which corresponded to the period when many leaf traits were relatively stable (Reich *et al.*, 1991). Gas exchange rates of individual leaves of each species were measured on sunny days between June 26 and July 25, 1998 (13 days), and June 12–24, July 9–24, 1999 (14 days). Rates in the 2 yr were determined at or near light-saturating conditions (mean PAR \pm SE: 1849 \pm 18 and 1616 \pm 9 $\mu\text{mol m}^{-2} \text{s}^{-1}$, in 1998 and 1999, respectively), at 25.5 \pm 177 \pm 0.1 and 26.6 \pm 177 \pm 0.1 °C (1998 and 1999, respectively), near ambient humidity (mean chamber VPD \pm SE: 1.65 \pm 177 \pm 0.03 and 1.51 \pm 177 \pm 0.02 kPa, in 1998 and 1999, respectively), and approximately at the CO₂ concentrations under which the plants were grown (356 \pm 1 or 549 \pm 1 \pm 109 $\mu\text{mol mol}^{-1}$). Hereafter, 368 and 560 $\mu\text{mol mol}^{-1}$ will be used to describe both measurement and growth CO₂ concentrations for clarity. Photosynthetic rates were calculated on both leaf area (A , $\mu\text{mol m}^{-2} \text{s}^{-1}$) and leaf mass (A_m , $\mu\text{mol g}^{-1} \text{s}^{-1}$) bases, and will be presented on an area basis unless otherwise denoted. Comparisons made between gas exchange measurements taken at growth CO₂ concentrations are referred to as ‘long-term responses’ ($A_{@growth}$, A_{560} vs A_{368}). The photosynthetic rate of each sample leaf of plants grown under ambient CO₂ was also determined at 560 $\mu\text{mol mol}^{-1}$ CO₂ concentration to assess the response of species to instantaneous CO₂ enrichment, referred to as ‘short-term responses’ (A'_{560} vs A_{368} , with A'_{560} referring to ambient grown plants measured at 560). This also allowed the comparison of rates of ambient and elevated CO₂ grown plants at a common CO₂ concentration ($A_{@560}$, A'_{560} vs A_{560}). Each of two replicate plots representing each of the four CO₂ \times N treatment combinations were sampled four times each year resulting in $n = 64$ for each species ($n = 32$ for *Lespedeza*, *Amorpha* and *Petalostemum* measured in one year only). The four samplings of each plot

were measured on separate days, at random time points, and were averaged for each replicate plot to incorporate day to day and within day variation in environmental conditions.

The projected areas of leaves used in gas exchange measurements were determined using a digital image analysis program (WinRhizo 3.9, Regent Instruments, Quebec, Canada). Leaves were then oven-dried (65 °C) to determine dry mass and specific leaf area (SLA, $\text{cm}^{-2} \text{g}^{-1}$). Each leaf used in gas exchange measurements was ground and analysed for tissue nitrogen concentration with an NA1500 C-N Analyser (Carlo-Erba Instruments, Milan, Italy). Intrinsic instantaneous water-use efficiency ($A_{@growth}/g_s$) and photosynthetic N-use efficiency (PNUE, $A_{@growth}/N_{area}$) were derived from gas exchange and tissue N data.

Data analysis

Data from four individual leaves, each measured on a separate day, were averaged for each replicate plot to provide an estimate of plot response to the treatment combinations within each year. Therefore, plot was the experimental unit used in ANOVA. For each of the nine variables analysed (Table 2), data from species measured in both 1998 and 1999 were analysed by repeated measures ANOVA including year in the model. Year as a main effect was significant ($P < 0.01$) for seven of the nine variables. The CO₂ treatment \times year interaction was not significant ($P > 0.25$) for any variable, however, in three cases the N treatment \times year interaction was significant ($P < 0.05$). Because the effect of year was often significant, year \times treatment interactions occasionally significant, and most importantly because some species were not measured in both years, we analysed and present statistics and data separately for each year. ANOVA was used in two complementary ways, including either species or functional group as a treatment effect for the following response variables: A , SLA, g_s , $A_{@growth}/g_s$, leaf N concentration (%N) and content (N_{area} , g N m^{-2}), and PNUE. In ANOVA, all treatment effects were considered fixed. Using F -tests, the effect of CO₂ (1 df) was tested against the random effect of ring nested within CO₂ (4 df). The main effect of functional group (3 df) was tested against the random effect of species nested within functional group (7 or 8 df). The main effects of species (10 or 11 df) and N (1 df), and all interactions, were tested against the residual error. To test the effect of CO₂ on the relationships between $A_{@560}-g_s$ and $A_{m,@560}-\%N$ we examined all data for individual species using ‘separate slopes’ analysis of covariance. We tested whether the slopes of regression lines varied among CO₂ treatments. If they did not differ significantly, the interaction term was removed from the model and ‘same slopes’ analyses were used to test for equality of intercepts between the CO₂ treatment regression lines. All analyses were conducted with statistical analysis software (JMP Version 3.16, SAS Institute Inc., Cary, NC, USA).

Table 1 Interspecific variation in proportional change in area-based net photosynthetic rate in response to elevated compared to ambient CO₂. Mean long-term responses are shown as the ratio of photosynthetic rates determined at growth CO₂ concentrations of 560 vs 368 μmol mol⁻¹ (A_{560}/A_{368}). Short-term responses to elevated CO₂ are shown as the ratio of photosynthetic rates of ambient CO₂ grown plants measured at 560 vs 368 μmol mol⁻¹ CO₂ (A'_{560}/A_{368}).

Species	1998				1999			
	Long-term Response (A_{560}/A_{368})		Short-term Response (A'_{560}/A_{368})		Long-term Response (A_{560}/A_{368})		Short-term Response (A'_{560}/A_{368})	
	High N	Low N	High N	Low N	High N	Low N	Low N	High N
Forbs								
<i>Achillea</i>	1.0	1.7	1.5	1.6	0.9	1.1	1.5	1.5
<i>Anemone</i>	0.9	1.1	1.5	1.6	0.8	0.8	1.5	1.6
<i>Solidago</i>	0.9	0.9	1.4	1.7	1.1	1.2	1.4	1.4
Legumes								
<i>Lupinus</i>	1.0	0.8	2.1	1.7	1.1	0.9	– ¹	–
<i>Lespedeza</i>	1.1	1.5	1.6	1.5	–	–	–	–
<i>Amorpha</i>	–	–	–	–	0.1	1.0	1.5	1.4
<i>Petalostemum</i>	–	–	–	–	1.1	0.9	1.4	1.3
C₃ Grasses								
<i>Agropyron</i>	1.1	1.1	1.7	1.5	1.4	0.8	1.5	1.4
<i>Bromus</i>	1.3	1.2	1.6	1.6	1.1	1.1	1.4	1.5
<i>Koeleria</i>	1.4	1.3	1.8	1.7	0.8	0.8	1.5	1.5
C₄ Grasses								
<i>Andropogon</i>	0.6	1.2	1.1	1.0	0.9	0.9	1.2	1.2
<i>Schizachyrium</i>	0.9	0.9	1.1	1.1	1.2	1.2	1.1	1.2
<i>Sorghastrum</i>	1.3	0.9	1.2	1.2	1.4	1.0	1.1	1.0

¹Data not available. A value of 1.0 indicates no difference in rates between CO₂ treatments. Statistics for long-term responses are shown in Table 2. The main effect of measurement CO₂ concentration for the short-term response $P > 0.0001$ in both years and in 1998: N effect $P = 0.99$, species \times N $P = 0.14$ 1999: N effect $P = 0.05$, species \times N = 0.01, in neither year was measurement CO₂ \times N significant.

Results

Net photosynthesis

Mean rates of light-saturated leaf net photosynthesis (A , μmol m⁻² s⁻¹) of the C₃ species grown at ambient CO₂ (368 μmol mol⁻¹) were increased on average by 64% and 44% in 1998 and 1999, respectively, after short-term exposure to elevated CO₂ (560 μmol mol⁻¹) (A'_{560} vs A_{368} , Table 1). By contrast, photosynthetic rates of C₃ species grown and measured at elevated CO₂ ('long-term exposure') were similar to or only slightly higher (mean difference +13% and +8% in 1998 and 1999, respectively) than plants grown and measured at ambient CO₂ (A_{560} vs A_{368}), and these differences were not statistically significant (Tables 1, 2; Fig. 1). Similarly, short-term exposure to 560 μmol mol⁻¹ of C₄ species grown under ambient CO₂ resulted in a 13% increase on average across years, while C₄ plants grown and measured at elevated CO₂ had similar rates of photosynthesis (mean difference –5% and +1% in 1998 and 1999, respectively) as those grown and measured at ambient CO₂ (Tables 1, 2; Fig. 1).

Hence, it appears that marked photosynthetic acclimation to elevated CO₂ occurred in all species in both years. When comparing the very modest proportional change in photosynthetic rates of elevated compared with ambient CO₂ grown

plants ('long-term response' A_{560}/A_{368}), with that of the substantial increase in proportional change in response to short-term increases in CO₂ (A'_{560}/A_{368}), pronounced acclimation of photosynthesis to growth under elevated CO₂ was revealed (Table 1). Moreover, comparing elevated and ambient CO₂-grown plants at a common measurement CO₂ concentration (A_{560} and A'_{560} , respectively), acclimation of photosynthesis was indicated by generally lower rates (–27% and –21% in 1998 and 1999, respectively) of elevated grown plants compared with ambient grown counterparts ($A_{@560}$; Table 2, Fig. 2).

In response to soil N treatments in 1998, plants grown under high N had 8% ($P = 0.08$) and 12% ($P = 0.008$) higher photosynthetic rates, on leaf area and mass bases, respectively (Table 2, Fig. 1a). However, not all species responded positively to the high N treatment (Table 2, species \times N interaction $P = 0.08$). *Lupinus*, an N-fixing legume, showed an average 34% decrease in $A_{@growth}$ under high N compared with low N (Fig. 1a). During 1999, overall N effects diminished such that $A_{@growth}$ was only slightly to moderately enhanced (+2% on average) in plants grown under high N with species responding similarly (Table 2, Fig. 1b). Although across most species high N increased rates of photosynthesis at least in the first year, the high N treatment had no significant effect on either short- or long-term photosynthetic responses to elevated CO₂. In fact, CO₂ \times N interactions were not significant for any response variable analysed (Table 1, 2).

Table 2 ANOVA probabilities ($P > F$) for treatment (CO_2 , N, and species) and main effects, and interactions, on leaf-level traits of grassland species (11 in 1998, 12 in 1999) grown at ambient ($368 \mu\text{mol mol}^{-1}$) and elevated ($560 \mu\text{mol mol}^{-1}$) CO_2 concentrations and low N (unamended soil) and high N treatments (addition of $4 \text{ g N m}^{-2} \text{ y}^{-1}$) over two growing seasons

Main effects & interactions	1998	$P > F$	A_{growth} ($\mu\text{mol m}^{-2} \text{ s}^{-1}$)	$A_{\text{m,growth}}$ ($\text{nmol g}^{-1} \text{ s}^{-1}$)	A_{560} ($\mu\text{mol m}^{-2} \text{ s}^{-1}$)	SLA ($\text{cm}^2 \text{ g}^{-1}$)	g_s ($\text{mmol m}^{-2} \text{ s}^{-1}$)	A_{growth}/g_s (mmol CO_2 $\text{mol H}_2\text{O}^{-1}$)	Leaf % N	Leaf N_{area} (gN m^{-2})	PNUE ($A_{\text{growth}}/N_{\text{area}}$) ($\mu\text{mol CO}_2 \text{ g N}^{-1} \text{ s}^{-1}$)
CO_2	1998	$P > F$	0.46 (+6) ¹	0.45 (+4)	0.03 (-28)	0.98 (4E)	0.05 (-23)	0.02 (+37)	0.04 (-12)	0.07 (-12)	0.18 (+13)
	1999	$P > F$	0.25 (+7)	0.66 (+3)	0.01 (-21)	0.37 (-2)	0.04 (-24)	0.01 (+43)	0.04 (-8)	0.21 (+8)	0.19 (+8)
N	1998	$P > F$	0.08 (+8)	0.008 (+12)	0.47 (+4)	0.55 (+1)	0.64 (-3)	0.0003 (+22)	0.0003 (+12)	0.003 (+13)	0.57 (-2)
	1999	$P > F$	0.44 (+2)	0.13 (+4)	0.50 (+2)	0.63 (+1)	0.86 (+1)	0.38 (+4)	0.75 (-1)	0.66 (-2)	0.60 (+3)
Species × CO_2	1998	$P > F$	0.21	0.48	0.07	0.004	0.93	0.04	0.007	0.49	0.02
Species × N	1999	$P > F$	0.85	0.45	0.26	0.02	0.82	0.38	0.25	0.56	0.56
	1998	$P > F$	0.08	0.0003	0.08	0.58	0.75	0.003	0.12	0.75	0.11
$\text{CO}_2 \times \text{N}$	1999	$P > F$	0.24	0.04	0.37	0.07	0.25	0.78	0.82	0.95	0.18
	1998	$P > F$	0.30	0.14	0.28	0.88	0.89	0.09	0.35	0.56	0.91
$\text{CO}_2 \times \text{N} \times \text{species}$	1999	$P > F$	0.10	0.08	0.29	0.75	0.81	0.72	0.98	0.54	0.17
	1998	$P > F$	0.19	0.02	0.47	0.14	0.28	0.31	0.99	0.99	0.19
	1999	$P > F$	0.57	0.52	0.74	0.63	0.65	0.75	0.02	0.09	0.12

¹Percent differences for main effects are shown below P -value. $P < 0.10$ are bold-faced. Species main effect $P < 0.0001$ for all response variables in both seasons. A_{growth} (area-based photosynthesis at growth CO_2 concentration); A_{m} (mass-based photosynthesis at growth CO_2 concentration); A_{560} (area-based photosynthesis measured at common CO_2 concentration of $560 \text{ mmol mol}^{-1}$); SLA (specific leaf area); g_s (stomatal conductance at the growth CO_2 concentration); A_{growth}/g_s (intrinsic instantaneous water-use efficiency); PNUE (photosynthetic N-use efficiency).




Fig. 1 Area-based rates of leaf net photosynthesis ($A_{@growth}$, $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) of 13 grassland species grown and measured under ambient (Open bars, $368 \mu\text{mol mol}^{-1}$) and elevated (hatched bars, $560 \mu\text{mol mol}^{-1}$) concentrations of CO_2 and low (unamended) and high (amended with $4 \text{ g N m}^{-2} \text{ y}^{-1}$) soil N. (a) 1998 (11 species) and (b) 1999 (12 species). Shown are least square means (\pm SE) from species \times $\text{CO}_2 \times$ N interaction in ANOVA. Species are arranged by functional group. See Table 2 for statistics.




Fig. 2 Area-based rates of net photosynthesis ($A_{@560}$, $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) of 13 grassland species measured at a common CO_2 concentration of $560 \mu\text{mol mol}^{-1}$. Plants were grown under ambient (open bars, $368 \mu\text{mol mol}^{-1}$) and elevated (hatched bars, $560 \mu\text{mol mol}^{-1}$) concentrations of CO_2 and low (unamended) and high (amended with $4 \text{ g N m}^{-2} \text{ y}^{-1}$) soil N. (a) 1998 (11 species) and (b) 1999 (12 species). Shown are least square means (\pm SE) from species \times $\text{CO}_2 \times$ N interaction in ANOVA. Species are arranged by functional group. See Table 2 for statistics.




Fig. 3 Leaf stomatal conductance to water vapor (g_s , $\text{mmol H}_2\text{O m}^{-2} \text{s}^{-1}$) of 13 grassland species grown and measured under ambient (open bars, $368 \mu\text{mol mol}^{-1}$) and elevated (hatched bars, $560 \mu\text{mol mol}^{-1}$) concentrations of CO_2 and low (unamended) and high (amended with $4 \text{ g N m}^{-2} \text{y}^{-1}$) soil N. Shown are least square means (\pm SE) from species \times $\text{CO}_2 \times$ N interaction in ANOVA. Species are arranged by functional group. See Table 2 for statistics.

Stomatal conductance and intrinsic instantaneous water-use efficiency

Stomatal conductance to water vapor (g_s) was on average 24% lower across all species and in both 1998 and 1999 in plants grown and measured at elevated compared with ambient CO_2 ($P = 0.05$, Table 2, Fig. 3). The reduced water loss, coupled with slightly enhanced $A_{\text{@growth}}$ in elevated CO_2 grown plants, resulted in an average 40% increase ($P = 0.02$) in intrinsic instantaneous water-use efficiency across all species ($A_{\text{@growth}}/g_s$, $\text{mmol CO}_2 \text{ mol H}_2\text{O}^{-1}$, Table 2).

High N did not significantly affect stomatal conductance, but in 1998 where high N enhanced net photosynthesis, $A_{\text{@growth}}/g_s$ was also increased (+22%, $P = 0.0003$), resulting in one of the few marginally significant interactions between CO_2 and N effects ($P = 0.09$, Table 2). In this case, the response of $A_{\text{@growth}}/g_s$ to elevated CO_2 was greater in magnitude under high N (43%) than under low N (23%) (data not shown).

Specific leaf area, leaf nitrogen and photosynthetic nitrogen-use-efficiency

Overall changes in SLA in response to either elevated CO_2 or added N were similar in both years but varied by species

(Table 3). Most species showed no difference or slight decreases in SLA between elevated and ambient CO_2 -grown plants. Most notable deviations from this pattern occurred in *Andropogon* in 1998 and *Lupinus* in 1999, in which SLA increased under elevated CO_2 by 26% and 17%, respectively, averaged across N treatments. Responses of these two species account for the observed significant species \times CO_2 interactions (Table 2).

The effects of CO_2 and N on leaf N concentration (%N) and content (N_{area} , g N m^{-2}) were also independent (Table 2). Across species and years, leaf N content declined by roughly 10% in elevated CO_2 grown plants (Table 2, Fig. 4). In 1998, leaf N content increased an average of 13% in leaves of plants grown at high N ($P = 0.003$, Table 2, Fig. 4). However in 1999, leaf N content of plants grown at high N did not differ from their low N grown counterparts ($P = 0.66$, Table 2, Fig. 4).

The effect of elevated CO_2 on leaf N content, coupled with its effect on photosynthesis, resulted in an 11% higher photosynthetic nitrogen-use-efficiency (PNUE, $A_{\text{@growth}}/N_{\text{area}}$, $\mu\text{mol CO}_2 \text{ g N}^{-1}$) on average across species (Table 2). In the first growing season, species PNUE responses to CO_2 enrichment ranged between a 21% decrease to a 59% increase in PNUE (species \times CO_2 interaction, $P = 0.02$, Table 2).

Functional group	Species	[CO ₂]	N Treatment			
			1998		1999	
			Low N	High N	Low N	High N
Forbs	<i>Achillea</i>	368	111.1	105.0	120.4	112.4
		560	105.7	102.2	106.4	101.5
	<i>Anemone</i>	368	178.8	161.1	149.1	167.2
		560	158.5	165.9	145.4	124.9
	<i>Solidago</i>	368	105.7	109.4	105.2	106.3
		560	104.2	108.1	104.8	113.8
Legumes	<i>Lupinus</i>	368	183.3	177.3	193.7	176.7
		560	166.9	189.4	232.1	200.6
	<i>Lespedeza</i>	368	159.7	146.4	– ²	–
		560	159.9	141.2	–	–
	<i>Amorpha</i>	368	–	–	153.1	169.4
		560	–	–	149.3	145.3
<i>Petalostemum</i>	368	–	–	114.9	112.5	
	560	–	–	95.5	108.0	
C ₃ Grasses	<i>Agropyron</i>	368	195.2	203.0	175.8	187.7
		560	195.9	182.0	164.9	188.4
	<i>Bromus</i>	368	198.6	203.3	166.3	163.0
		560	178.8	223.4	166.7	162.4
	<i>Koeleria</i>	368	138.5	128.5	89.8	102.5
		560	112.8	117.9	86.5	93.5
C ₄ Grasses	<i>Andropogon</i>	368	225.2	256.7	181.1	189.0
		560	323.2	285.8	185.8	197.3
	<i>Schizachyrium</i>	368	210.3	217.9	204.2	185.7
		560	193.4	221.5	194.6	184.5
	<i>Sorghastrum</i>	368	191.4	207.1	168.1	173.5
		560	195.5	186.7	157.5	174.7

Table 3 Specific leaf area (SLA, cm² g⁻¹) of grassland species (11 in 1998, 12 in 1999) grown under 368 and 560 μmol mol⁻¹ CO₂ and low and high soil N treatments¹

¹Presented are least square means (± 12.0 cm² g⁻¹ (1998), ± 8.8 cm² g⁻¹ (1999)) from species × CO₂ × N interaction in ANOVA. See Table 2 for statistics. ²Data not available.

A_{@560}-g_s and A_{m,@560}-%N relationships

We examined the relationships between A_{@560}-g_s (Fig. 5a,b) and A_{m,@560}-%N (Fig. 5c,d) across species to evaluate possible mechanisms to explain the photosynthetic acclimation to growth under elevated CO₂. As these associations varied among species, four specific cases were chosen that most clearly show the two types of relationships found. Some species grown under elevated CO₂ responded with decreases in photosynthesis that were proportional to decreases in stomatal conductance (Fig. 5a). However in other species, photosynthesis was lower at a given stomatal conductance in elevated compared with ambient CO₂ grown plants (Fig. 5b). Qualitatively, roughly half of the 13 species, in at least one of the 2 yr, showed a relationship between A_{@560} and g_s closer to the former. Declines in photosynthesis of elevated CO₂ grown plants were associated with decreased N concentration in six of the 13 species (Fig. 5c), while the others had lower photosynthetic rates at a given leaf N in elevated compared with ambient CO₂ grown plants (Fig. 5d). Whether species demonstrated one type of relationship vs the other was not related to their functional groupings.

Functional groups

For analyses on data pooled across species within their functional groupings, statistically significant interactions between functional groups and CO₂ or N treatments were few and occurred predominately in 1998 (Fig. 6a,b). In addition, CO₂ and soil N treatment interactions were nonsignificant in all cases. The enhancements in leaf-level photosynthetic rates by growth in elevated CO₂ of the C₃ species (C₃ grasses, legumes, and nonleguminous forbs) were greater (average +10%) than that of the C₄ species (average +1%) across both seasons. The most positive responses were among the C₃ grasses, which had 21% and 11% higher rates of photosynthesis when grown under elevated compared with ambient CO₂ in 1998 and 1999, respectively (Fig. 6a,c).

In terms of leaf N content, there was an interaction between functional groups and CO₂ in 1998 but not 1999. (Fig. 6a,c; *P* = 0.02). The effect of CO₂ concentration on PNUE was also less in the C₄ grasses than the other functional groups in both 1998 and 1999, but this interaction was significant in 1998 only (Fig. 6a, *P* = 0.01). In 1998, the responses of A, A_{@growth}/g_s, and leaf N content to the high




Fig. 4 Area-based leaf N content (N_{area} g N m^{-2}) of 13 grassland species grown and measured under ambient (open bars, $368 \mu\text{mol mol}^{-1}$) and elevated (hatched bars, $560 \mu\text{mol mol}^{-1}$) concentrations of CO_2 and low (unamended) and high (amended with $4 \text{ g N m}^{-2} \text{ y}^{-1}$) soil N. (a) 1998 (11 species) and (b) 1999 (12 species). Shown are least square means (\pm SE) from species \times $\text{CO}_2 \times$ N interaction in ANOVA. Species are arranged by functional group. Each year was analysed separately. See Table 2 for statistics.




Fig. 5 Relationships between leaf net photosynthesis (A_{560}/A_{360}) and leaf stomatal conductance (g_{s560} , a,b) or leaf N concentration (% , c, d) of foliage grown under ambient (closed circles, $368 \mu\text{mol mol}^{-1}$) and elevated (open circles, $560 \mu\text{mol mol}^{-1}$) concentrations of CO_2 and measured at a common CO_2 concentration of $560 \mu\text{mol mol}^{-1}$. These species were chosen to represent the variation in response across the 13 grassland species included in this study. Shown are data from individual measurements. Correlation coefficients (r^2) for A_{560}/A_{360} vs g_s : (a) *Amorpha* 99 ($r^2 = 0.75$, $P < 0.0001$) (b) *Bromus* 99 (ambient CO_2 $r^2 = 0.39$, $P = 0.01$; elevated CO_2 $r^2 = 0.41$, $P = 0.007$); for A_{560}/A_{360} vs %N: (c) *Lupinus* 98 ($r^2 = 0.46$, $P = 0.0004$) (d) *Bromus* 98 (ambient CO_2 $r^2 = 0.47$, $P = 0.003$, elevated CO_2 $r^2 = 0.79$, $P < 0.0001$). See text for further explanation of analysis.

soil N treatment were positive in each functional group but the legumes responded in the opposite direction ($\text{fx}_{\text{grp}} \times \text{N}$ interactions, Fig. 6b). The few cases in which species pooled by functional groups showed differential responses to CO_2 concentration or soil N treatments were due to either the legumes or the C_4 grasses.

Discussion

Acclimation of photosynthesis to elevated CO_2 concentration and increased soil N

Photosynthetic rates are well known to increase in C_3 plants in response to short-term increases in CO_2 concentrations (Bazzaz, 1990; Drake *et al.*, 1997). In this study, photosynthetic rates increased an average of 55% for the C_3 species and 13% for the C_4 species in response to short-term CO_2 enrichment, which is comparable in magnitude with the 60% average increase found in other studies of C_3 species (Strain & Cure, 1994; Curtis, 1996; Drake *et al.*, 1997). Theory based on simple CO_2 diffusion into C_3 leaves suggests that a photosynthetic enhancement of 55% is expected with $+200 \mu\text{mol mol}^{-1}$ enrichment in CO_2 (as in this study) and no photosynthetic acclimation (Katul *et al.*, 2000).




Fig. 6 Relative percent changes in response traits by functional groupings ((elevated – ambient)/ambient × 100). F, forbs; L, legumes; C₃, C₃ grasses; C₄, C₄ grasses; (a) and (b) 1998 (c) and (d) 1999; (a) and (c) CO₂ treatment effects pooled across N treatments (black bars) (b) and (d) N treatment effects pooled across CO₂ treatment (hatched bars). Significant functional group × CO₂ treatment (f_{xgrp} × CO₂) and functional group × N treatment (f_{xgrp} × N) interactions are given. No CO₂ × N interactions were significant. A_{@growth} (area-based photosynthesis, μmol m⁻² s⁻¹), A_m (mass-based photosynthesis, nmol g⁻¹ s⁻¹), specific leaf area (SLA, cm² g⁻¹), g_s (stomatal conductance, mmol m⁻² s⁻¹), Leaf N_{area} (gN m⁻²), A_{@growth}/g_s (intrinsic instantaneous water-use efficiency, mmol CO₂ mol H₂O⁻¹), PNUE (photosynthetic N-use efficiency, μmol CO₂ g N⁻¹ s⁻¹).

The degree of relative photosynthetic responsiveness to elevated CO₂ over longer-terms varies substantially. The long-term enhancement of photosynthesis by elevated CO₂ we observed was modest and not statistically significant in either year. C₃ species on average increased photosynthesis when grown under elevated CO₂ by 13% and 8%, in 1998 and 1999, respectively, and C₄ species demonstrated negligible responses in both years. This is in contrast to many field-based studies that have found strong and persistent stimulation of photosynthetic rates in C₃ species grown under elevated CO₂ over one to three growing seasons (Ellsworth *et al.*, 1995; Jackson *et al.*, 1995; Drake *et al.*, 1996; Stirling *et al.*, 1997; Bryant *et al.*, 1998; Curtis *et al.*, 2000) as well as positive enhancements for C₄ species (Read *et al.*, 1997; Wand *et al.*, 1999, but not Tissue *et al.*, 1995). Recent reviews summarize long-term photosynthetic enhancements in response to growth

under elevated CO₂ on the average of 30% and 24% in C₃ and C₄ Poaceae species, respectively (Wand *et al.*, 1999), and 50% and 51% across 41 woody C₃ species (Curtis, 1996), and 15 field-based on forest tree species (Medlyn *et al.*, 1999), respectively. However, other studies demonstrate limited stimulation of photosynthesis of plants grown for weeks to months under elevated CO₂ (Tissue & Oechel, 1987; Li *et al.*, 1999; Roumet *et al.*, 2000) or a loss of the initial stimulation in photosynthesis over time (Oechel *et al.*, 1994; Körner *et al.*, 1997).

Species variation in responsiveness of photosynthesis to elevated CO₂ can be explained by differences in the extent of photosynthetic acclimation in plants grown under elevated CO₂. In our study on nutrient poor soil, species on average demonstrated 80% acclimation of photosynthesis to elevated atmospheric CO₂ concentrations as judged by comparing the

magnitude of long-term photosynthetic enhancements with elevated CO₂ (A_{560} vs A_{368}) to the photosynthetic response to short-term exposure to elevated CO₂ (A'_{560} vs A'_{368}). This is much larger than shown in most previous studies (Curtis, 1996; Medlyn *et al.*, 1999; Wand *et al.*, 1999, but not Tissue & Oechel, 1987).

5

Is photosynthetic acclimation to elevated CO₂ modulated by soil N supply? The magnitude of the CO₂ effect on gas exchange and other leaf traits in these 13 grassland species was apparently independent of soil N supply in these two years. In only three specific cases (*Achillea* 98 and 99, and *Lespedeza* 98, Fig. 1) was there an increase in magnitude of enhancement of photosynthesis due to elevated CO₂ under high N, though not statistically significant (Table 2). Several other studies have found plants to be comparably responsive to CO₂ at both high and low nutrient concentrations (Hättenschwiler & Körner, 1996; Lloyd & Farquhar, 1996; Körner *et al.*, 1997; Cotrufo *et al.*, 1998). However, many studies have found greater photosynthetic responsiveness to elevated CO₂ at higher N availability (Curtis, 1996; Miglietta *et al.*, 1996; Rogers *et al.*, 1998; Sims *et al.*, 1998; Wolfe *et al.*, 1998; Weerakoon *et al.*, 1999; Curtis *et al.*, 2000). Moreover, in some cases, there is evidence of acclimation of photosynthesis under CO₂ enrichment only under certain conditions such as low nutrient supply (Jones *et al.*, 1996; Miglietta *et al.*, 1996; Rogers *et al.*, 1998; Sims *et al.*, 1998; Weerakoon *et al.*, 1999). In our study, the positive effects of high N on traits such as net photosynthetic rates, $A_{@growth}/g_s$, and leaf N concentrations were found only in the first growing season and diminished by the second year. While this difference in response between two years does not necessarily constitute a trend, it was contrary to expectations that the effects of adding N into the system each year would result in stronger N effects over time. This minimal N effect may have contributed to the lack of a CO₂ × N interaction.

6

One of the most consistent responses across species and growing seasons in our study was a decline in stomatal conductance to water vapor (g_s). All species decreased g_s by an average 24% when grown under elevated CO₂. While some studies report little or inconsistent effects of elevated CO₂ on g_s (Gunderson & Wullschleger, 1994; Ellsworth *et al.*, 1995; Curtis, 1996; Stirling *et al.*, 1997) the more common response is a decline in g comparable in magnitude to this study (Roumet *et al.*, 2000). Reviews cite average declines in g_s of 24% and 29% in C₃ and C₄ Poaceae species, respectively (Wand *et al.*, 1999), 23% across 23 tree species (Field *et al.*, 1995), and 34% across crop species (Kimball & Idso, 1983) in elevated compared with ambient CO₂ grown plants.

As leaf N concentration declined an average of 13%, carbon assimilation expressed per unit leaf N (i.e. PNUE) increased an average of 11% across the species in this study. However, this response was not statistically significant and not consistent across all species. Cotrufo *et al.* (1998) found a comparable reduction in tissue N concentrations (average

of 14%) of elevated CO₂ grown C₃ and C₄ plants across 75 published studies and other studies have also found that PNUE increases in response to growth under elevated CO₂ (Bryant *et al.*, 1998; Tjoelker *et al.*, 1998; Peterson *et al.*, 1999b; Curtis *et al.*, 2000), however, in some cases this did not occur consistently (Roumet *et al.*, 2000).

Several possible explanations for the acclimation of photosynthesis in plants grown under elevated CO₂ over longer terms have been proposed. Hypotheses include possible stomatal limitations of photosynthesis due to reduced g_s (Drake *et al.*, 1997), or nonstomatal limitations such as reduced tissue N concentrations potentially leading to a reduced photosynthetic capacity in plants grown under elevated CO₂ (Peterson *et al.*, 1999), or a potential feedback inhibition of photosynthesis induced by an accumulation of excess carbohydrates (Farrar & Williams, 1991; Stitt, 1991).

We examined the relationships between $A_{@560}-g_s$ and $A_{m,@560}-\%N$ to test the first two hypotheses. Results varied among species, but did provide evidence for both possible mechanisms. Some species grown under elevated CO₂ responded with decreases in photosynthesis that were proportional to decreases in g_s (Fig. 5a). In these cases, there was a concomitant decrease in C_i : C_a (data not shown) further suggesting that the decline in g_s of elevated CO₂ grown plants was associated with a lower intercellular CO₂ supply and reduced photosynthetic rates. In other species, photosynthesis was lower at a given g_s , with similar or slightly higher C_i : C_a, in elevated compared with ambient CO₂ grown plants (Fig. 5b), suggests that nonstomatal limitations are also involved.

A potential nonstomatal limitation involves the commonly strong relationship between tissue N and photosynthesis. Regarding the N hypothesis to explain photosynthetic acclimation, we did find a decline in photosynthesis in elevated compared with ambient CO₂ grown plants in proportion to the change in leaf N concentration (Fig. 5c) in roughly half the species. Thus CO₂-induced decreases in leaf N concentration is associated with reduced photosynthetic potential probably via changes in N-rich photosynthetic enzymes that are reflected in total leaf N. However, in other cases, photosynthesis was lower at a given leaf N in elevated compared with ambient CO₂ grown plants (Fig. 5d) suggesting that stomatal or other nonstomatal limitations explain the observed acclimation in such cases.

Studies have found that an increase in total nonstructural carbohydrates (TNC) correlates with decreased photosynthesis in elevated CO₂ grown plants (Tjoelker *et al.*, 1998; Roumet *et al.*, 2000), however, others report increased TNC without a decrease in photosynthetic enhancement (Wullschleger *et al.*, 1992; Will & Ceulemans, 1997). Total nonstructural carbohydrates determined from leaves collected from the same plots used for gas exchange in this study in 1999, indicate an overall 24% greater TNC concentration in elevated CO₂ grown foliage, but this varies substantially across species (M. G. Tjoelker *et al.*, unpublished). Because acclimation of

photosynthesis occurs across all the species in the study, but not all species increased TNC when grown under elevated compared with ambient CO₂, these data are not sufficient to support or refute this as a possible explanation of the photosynthetic acclimation seen in this study.

Most species showed intermediate responses with respect to the examples shown in Fig. 5. Considering this evidence of stomatal and nonstomatal limitations on photosynthesis, no single mechanism explains the magnitude of photosynthetic acclimation seen across all the species in this study, and the data suggests that even within a single species a combination of these mechanisms are possible. Thus, we can ascribe the low or negligible photosynthetic enhancement observed at our site to the combined effects of decreased g_s , decreased leaf N concentration, and possibly increased TNC in response to growth under elevated CO₂.

Functional groups

In our study, the grouping of species into the functional classifications: C₃ grasses, C₄ grasses, legumes, and nonleguminous forbs, are based on discrete physiological and growth form traits commonly used to group species. Our objective was to evaluate whether these categories are helpful in explaining variation in species response to elevated CO₂ and increased N supply across the 13 perennial prairie species in this study. The C₄ grasses are grouped owing to their photosynthetic pathway that concentrates CO₂ in bundle sheath cells, which effectively increases the concentration of CO₂ at the site of carboxylation and therefore, presumably results in or near saturation at current CO₂ levels. As predicted, the C₄ species were less responsive than the C₃ species to elevated CO₂ over the long-term with negligible effects on photosynthesis compared with a 10% average increase for all the C₃ species combined. While the legumes, unique in the ability to symbiotically fix N₂, responded to the high N treatment oppositely, in terms of net photosynthesis, A/g_s, and tissue N, relative to the other functional groups. However, species pooled by functional groups responded differently to elevated CO₂ or high soil N treatments in relatively few cases, only for the C₄ grasses and the legumes, and predominantly in the first year of growth under CO₂ and N treatments. In a companion study by Reich *et al.* (2001), which looked at plot level traits such as total biomass, total plant N, soil solution N and soil water on these same monoculture plots, it was also found that in general, C₄ grasses were less responsive to elevated CO₂ than all C₃ species as a group and legumes were less responsive to the N treatment. Overall, the variation in species responsiveness to elevated CO₂ and soil N supply was generally unrelated to their functional groupings.

Summary

For this diverse set of species with varying ecophysiology, elevated CO₂ only modestly stimulated photosynthesis over

the longer term, and not in all cases. However, rates of stomatal conductance declined consistently across species. In general, soil N supply did not modulate species responses to elevated CO₂. In addition, the variation in species response was not exclusively explained by their functional groupings suggesting that current functional classifications may not be sufficient for understanding leaf-level physiological responses to elevated CO₂ and N. The extent to which wild species acclimate photosynthesis to elevated CO₂, and how the efficiency of resource use is affected by growth under elevated CO₂, appear to be critical in determining plant growth responses to changing atmospheric CO₂ and N deposition.

Acknowledgements

We acknowledge funding from the US Department of Energy and the Cedar Creek Natural History Area Long-Term Ecological Research site in support of this research. We thank G. Hendrey for the original FACE approach and D. Bahauddin and K. Wrage for constructing and operating the FACE system. We are grateful to Ann Karpinski, Jenny Goth, Wendy Bengtson, and Doreen Hoover for assistance with gas exchange measurements and sample preparation. This study was also supported by NSF Pre-doctoral and University of Minnesota Graduate School Doctoral Dissertation Fellowship Awards to Tali D. Lee.

References

- Bazzaz FA. 1990. The response of natural ecosystems to the rising global CO₂ levels. *Annual Review of Ecology and Systematics* 21: 167–196.
- Bowes G. 1993. Facing the inevitable: plants and increasing atmospheric CO₂. *Annual Review of Plant Physiology and Plant Molecular Biology* 44: 309–332.
- Bryant J, Taylor G, Frehner M. 1998. Photosynthetic acclimation to elevated CO₂ is modified by source: sink balance in three component species of chalk grassland swards grown in a free air carbon dioxide enrichment (FACE) experiment. *Plant, Cell & Environment* 21: 159–168.
- Cotrufo MF, Ineson P, Scott A. 1998. Elevated CO₂ reduces the nitrogen concentration of plant tissues. *Global Change Biology* 4: 43–54.
- Curtis PS. 1996. A meta-analysis of leaf gas exchange and nitrogen in trees grown under elevated carbon dioxide. *Plant, Cell & Environment* 19: 127–137.
- Curtis PS, Vogel CS, Wang X, Pregitzer KS, Zak DR, Lussenhop J, Kubiske M, Teeri JA. 2000. Gas exchange leaf nitrogen, and growth efficiency of *Populus tremuloides* in a CO₂-enriched atmosphere. *Ecological Applications* 10: 3–17.
- Díaz S. 1995. Elevated CO₂ responsiveness, interactions at the community level and plant functional types. *Journal of Biogeography* 22: 289–295.
- Drake BG, González-Meler MA, Long SP. 1997. More efficient plants: a consequence of rising atmospheric CO₂? *Annual Review of Plant Physiology and Plant Molecular Biology* 48: 609–639.
- Drake BG, Muehe MS, Peresta G, González-Meler MA, Matamala. 1996. Acclimation of photosynthesis, respiration and ecosystem carbon flux of a wetland on Chesapeake Bay, Maryland to elevated atmospheric CO₂ concentration. *Plant and Soil* 187: 111–118.
- Ellsworth DS, Oren R, Huang C, Phillips N, Hendrey GR. 1995. Leaf and canopy responses to elevated CO₂ in a pine forest under free-air CO₂ enrichment. *Oecologia* 104: 139–146.

- Farrar JF, Williams ML. 1991. The effects of increased atmospheric carbon dioxide and temperature on carbon partitioning, source-sink relations and respiration. *Plant, Cell & Environment* 14: 819–830.
- Field CB, Jackson RB, Mooney HA. 1995. Stomatal responses to increased CO₂: implications from the plant to the global scale. *Plant, Cell & Environment* 18: 1214–1225.
- Gunderson CA, Wullschlegel SD. 1994. Photosynthetic acclimation in trees to rising atmospheric CO₂: a broader perspective. *Photosynthesis Research* 39: 369–388.
- Hättenschwiler S, Körner C. 1996. Effects of elevated CO₂ and increased nitrogen deposition on photosynthesis and growth of understory plants in spruce model ecosystems. *Oecologia* 106: 172–180.
- Idso KE, Idso SB. 1994. Plant responses to atmospheric CO₂ enrichment in the face of environmental constraints: a review of the past 10 years research. *Agricultural and Forest Meteorology* 69: 153–203.
- Jackson RB, Luo Y, Cardon ZG, Sala OE, Field CB, Mooney HA. 1995. Photosynthesis, growth and density for the dominant species in a CO₂-enriched grassland. *Journal of Biogeography* 22: 221–225.
- Jones MB, Jongen M, Garvey L, Baxter R. 1996. The physiological responses of a European grassland species to elevated CO₂. *Aspects of Applied Biology* 45: 155–161.
- Katul GG, Ellsworth DS, Lai C. 2000. Modeling assimilation and intercellular CO₂ from measured conductance: a synthesis of approaches. *Plant, Cell & Environment* 23: 1313–1328.
- Kimball BA, Idso SB. 1983. Increasing atmospheric CO₂: effects on crop yield, water use and climate. *Agricultural Water Management* 7: 55–72.
- Körner Ch, Diemer M, Schächli B, Miklaus P, Arnone III J. 1997. The responses of alpine grassland to four seasons of CO₂ enrichment: a synthesis. *Acta Oecologia* 18: 165–175.
- Larigauderie A, Hilbert DW, Oechel WC. 1988. Effect of CO₂ enrichment and nitrogen availability on resource acquisition and resource allocation in a grass, *Bromus mollis*. *Oecologia* 77: 544–549.
- Leadley PW, Körner Ch. 1996. Effects of elevated CO₂ on plant species dominance in a highly diverse calcareous grassland. In: Körner CH, Bazzaz FA, eds. *Carbon dioxide, populations, and communities*. San Diego, CA, USA: Academic Press, Inc., 159–175.
- Lewin KF, Hendrey GR, Nagy J, LaMorte R. 1994. Design and application of a free-air carbon dioxide enrichment facility. *Agricultural and Forest Meteorology* 70: 15–29.
- Li JH, Dijkstra P, Hinkle CR, Wheeler RM, Drake BG. 1999. Photosynthetic acclimation to elevated atmospheric CO₂ concentration in the Florida scrub-oak species *Quercus myrtifolia* growing in their native environment. *Tree Physiology* 19: 229–234.
- Lloyd J, Farquhar GD. 1996. The CO₂ dependence of photosynthesis, plant growth responses to elevated atmospheric CO₂ concentrations and their interaction with soil nutrient status. I. General principles and forest ecosystems. *Functional Ecology* 10: 4–32.
- McMurtie RE, Comins HN. 1996. The temporal response of forest ecosystems to doubled atmospheric CO₂ concentration. *Global Change Biology* 2: 49–57.
- Medlyn BE, Badeck F-W, de Pury DGG, Barton CVM, Broadmeadow M, Ceulemans R, deAngelis P, Forstreuter M, Jach ME, Kellomäki S, Laitat E, Marek M, Phillipot S, Rey A, Strassmeyer J, Laitinen K, Loizon R, Portier B, Roberntz P, Wang K, Jarvis PG. 1999. Effects of elevated [CO₂] on photosynthesis in European forest species: a meta-analysis of model parameters. *Plant, Cell & Environment* 22: 1475–1496.
- Miglietta F, Giuntoli A, Bindi M. 1996. The effect of free air carbon dioxide enrichment (FACE) and soil nitrogen availability on the photosynthetic capacity of wheat. *Photosynthesis Research* 47: 281–290.
- Nijs I, Behaeghe T, Impens I. 1995. Leaf nitrogen content as a predictor of photosynthetic capacity in ambient and global change conditions. *Journal of Biogeography* 22: 177–183.
- Oechel WC, Cowles S, Grulke N, Hastings SJ, Lawrence B, Prudhomme T, Riechers G, Strain B, Tissue D, Vourlitis G. 1994. Transient nature of CO₂ fertilization in Arctic tundra. *Nature* 371: 500–503.
- Peterson AG, Ball JT, Luo Y, Field CB, Reich PB, Curtis PB, Griffin KL, Gunderson CA, Norby RJ, Tissue DT, Forstreuter M, Rey A, Vogel CS, CMEAL participants. 1999. The photosynthesis-leaf nitrogen relationship at ambient and elevated atmospheric carbon dioxide: a meta-analysis. *Global Change Biology* 5: 331–346.
- Polley HW. 1997. Implications of rising atmospheric carbon dioxide concentration for rangelands. *Journal of Range Management* 50: 561–577.
- Poorter H. 1998. Do slow-growing species and nutrient-stressed plants respond relatively strongly to elevated CO₂? *Global Change Biology* 4: 693–697.
- Rastetter EB, Ågren GI, Shaver GR. 1997. Responses of N-limited ecosystems to increased CO₂: a balanced-nutrition, coupled-element-cycles model. *Ecological Applications* 7: 444–460.
- Reich PB, Knops J, Tilman D, Craine J, Ellsworth D, Tjoelker M, Lee T, Wedin D, Naeem S, Bahaeddin D, Hendrey G, Jose S, Wrage K, Goth J, Bengston W. 2001a. Plant diversity enhances ecosystem responses to elevated CO₂ and nitrogen deposition. *Nature* (In press.)
- Reich PB, Tilman D, Craine J, Ellsworth D, Tjoelker M, Knops J, Wedin D, Naeem S, Bahaeddin D, Goth J, Bengston W, Lee TD. 2001b. Do species and functional groups differ in acquisition and use of C, N and water under varying atmospheric CO₂ and N deposition regimes? A field-test with 16 grassland species. *New Phytologist* (In press.)
- Reich PB, Walters MB, Ellsworth DS. 1991. Leaf age and season influence the relationships between leaf nitrogen, leaf mass per area and photosynthesis in maple and oak trees. *Plant, Cell & Environment* 14: 251–259.
- Reich PB, Walters MB, Ellsworth DS. 1997. From tropics to tundra: Global convergence in plant functioning. *Proceedings of the National Academy of Sciences, USA* 94: 13730–13734.
- Rogers A, Fischer BU, Bryant J, Frehner M, Blum H, Raines CA, Long SP. 1998. Acclimation of photosynthesis to elevated CO₂ under low-nitrogen nutrition is affected by the capacity for assimilate utilization. Perennial ryegrass under free-air CO₂ enrichment. *Plant Physiology* 118: 683–689.
- Roumet C, Garnier E, Suzor H, Salager JL, Roy J. 2000. Short and long-term responses of whole-plant gas exchange to elevated CO₂ in four herbaceous species. *Environmental & Experimental Botany* 43: 155–169.
- Schimel DS. 1995. Terrestrial ecosystems and the carbon cycle. *Global Change Biology* 1: 77–91.
- Sims DA, Luo Y, Seeman JR. 1998. Comparison of photosynthetic acclimation to elevated CO₂ and limited nitrogen supply in soybean. *Plant, Cell & Environment* 21: 945–952.
- Stirling CM, Davey PA, Williams TG, Long SP. 1997. Acclimation of photosynthesis to elevated CO₂ and temperature in five British native species of contrasting functional type. *Global Change Biology* 3: 237–246.
- Stitt M. 1991. Rising CO₂ levels and their potential significance for carbon flow in photosynthetic cells. *Plant, Cell & Environment* 14: 741–762.
- Strain BR, Cure JD. 1994. *Direct effects of atmospheric CO₂ enrichment on plants and ecosystems: an updated bibliographic database*. Oak Ridge, TN, USA: Oak Ridge National Laboratory/CDIAC-70.
- Tilman D. 1987. Secondary succession and the pattern of plant dominance along experimental nitrogen gradients. *Ecological Monographs* 57: 189–214.
- Tissue DT, Oechel WC. 1987. Response of *Eriophorum vaginatum* to elevated CO₂ and temperature in the Alaskan tussock tundra. *Ecology* 68: 401–410.
- Tjoelker MG, Oleksyn J, Reich PB. 1998. Seedlings of five boreal species differ in acclimation of net photosynthesis to elevated CO₂ and temperature. *Tree Physiology* 18: 715–726.
- Vitousek PM, Aber JD, Howarth RW, Likens GE, Matson PA, Schindler DW, Schlesinger WH, Tilman DG. 1997. Human alteration of the global nitrogen cycle: sources and consequences. *Ecological Applications* 7: 737–750.
- Ward SE, Midgley GF, Jones MH, Curtis PS. 1999. Responses of wild

- C₄ and C₃ grasses (*Poaceae*) species to elevated atmospheric CO₂ concentration: a meta-analytic test of current theories and perceptions. *Global Change Biology* 5: 723–741.
- Wedin DA, Tilman D. 1996. Influence of nitrogen loading and species composition on the carbon balance of grasslands. *Science* 274: 1720–1723.
- Weerakoon WM, Olszyk DM, Moss DN. 1999. Effect of nitrogen nutrition on responses of rice seedlings to carbon dioxide. *Agriculture, Ecosystems and Environment* 72: 1–8.
- Will RE, Ceulemans R. 1997. Effects of elevated CO₂ concentration on photosynthesis, respiration and carbohydrate status of coppice *Populus* hybrids. *Physiologia Plantarum* 100: 933–939.
- Wolfe DW, Gifford RM, Hilbert D, Luo Y. 1998. Integration of photosynthetic acclimation to CO₂ at the whole-plant level. *Global Change Biology* 4: 879–893.
- Wullschlegel SD, Norby RJ, Hendrix DL. 1992. Carbon exchange rates, chlorophyll content and carbohydrate status of two forest tree species exposed to carbon dioxide enrichment. *Tree Physiology* 10: 21–31.

Author Query Form

Journal: New Phytologist

Article: 095

Dear Author,

During the preparation of your manuscript for publication, the questions listed below have arisen. Please attend to these matters and return this form with your proof.

Many thanks for your assistance.

Query Refs.	Query	Remarks
1	VPD – please give in full	
2	Read <i>et al.</i> 1997 has not been included in the list	
3	but not Tissue <i>et al.</i>, 1995 – this seems a little unclear. Is it correct?	
4	Tissue <i>et al.</i> 1995 has not been included in the list	
5	but not Tissue & Oechel, 1987 – again, this is a little unclear. Is the text correct?	
6	Miglietta 1996 has been changed to Miglietta <i>et al.</i> (1996) so that this citation matches the list	
7	While the legumes, unique in the ability to symbiotically fix N₂, responded to the high N treatment oppositely, in terms of net photosynthesis, A/g_s, and tissue N, relative to the other functional groups. – this sentence is somewhat unclear. Please clarify	
8	Please supply Matamala's initials	
9	Proceedings of the National Academy of Sciences, USA – 'USA' has been added. Is this ok?	